arXiv:1507.02749v1 [math.AT] 9 Jul 2015

Perfect Morse Function on $SO(n)$

Mehmet Solgun
Bilecik Seyh Edebali University, Faculty of Sciences and Arts
Department of Mathematics, Bilecik, TURKEY
mehmet.solgun@bilecik.edu.tr

July 13, 2015

Abstract

In this work, we define a Morse function on $SO(n)$ and show that this function is indeed a perfect Morse function.

Keywords: $SO(n)$, Morse functions, Perfect Morse functions. Mathematics Subject Classification 2010:57R70, 58E05

1 Introduction

The main point of Morse Theory, which was introduced in [6], is investigating the relation between shape of a smooth manifold M and critical points of a specific real-valued function $f : M \to \mathbb{R}$, that is called *Morse function.* [5] and [4] are two of main sources about this subject, so mostly we will use their beautiful tools for defining a Morse function on $SO(n)$. Also, we will refer [2] to use homological properties and to determine the Poincaré polynomial of $SO(n)$. Perfect Morse functions are widely studied in [7], that is one of our inspiration to show that the function, we defined, is also perfect.

2 Preliminaries

In this section, we give some definitions and theorems which will be used in this paper.

Definition 2.1 *Let M be an n-dimensional smooth manifold and $f : M \to \mathbb{R}$ be a smooth function. A point $p_0 \in M$ is said to be a critical point of M if we have*

$$\frac{\partial f}{\partial x_1} = 0,\ \frac{\partial f}{\partial x_2} = 0,\ ...,\ \frac{\partial f}{\partial x_n} = 0 \tag{2.1}$$

with respect to a coordinate system $\{x_1, x_2, ..., x_n\}$ around p_0.

A point $c \in \mathbb{R}$ is said to be *a critical value of* $f : M \to \mathbb{R}$, if $f(p_0) = c$ for a critical point p_0 of f.

Definition 2.2 *Let* p_0 *be a critical point of the function* $f : M \to \mathbb{R}$*. The Hessian of* f *at he point* p_0 *is the* $n \times n$ *matrix*

$$H_f(p_0) = \begin{bmatrix} \frac{\partial^2 f}{\partial {x_1}^2}(p_0) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(p_0) \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(p_0) & \cdots & \frac{\partial^2 f}{\partial {x_n}^2}(p_0) \end{bmatrix} \tag{2.2}$$

Since $\frac{\partial^2 f}{\partial x_i \partial x_j}(p_0) = \frac{\partial^2 f}{\partial x_j \partial x_i}(p_0)$, the Hessian of f is a symmetric matrix.

Let p_0 be a critical point of f and $c_0 \in \mathbb{R}$ such that $f(p_0) = c_0$. Then, c_0 is said to be *a critical value of* f. If p_0 is a regular point of f, then c_0 is said to be a *regular value of* f.

If a is a regular value of f, it can be shown that the set $f^{-1}(a) = \{p \in M | f(p) = a\}$ is an $n-1$ dimensional manifold [1].

Definition 2.3 *A critical point of a function* $f : M \to \mathbb{R}$ *is called "non-degenerate point of* f*" if* $det H_f(p_0) \neq 0$*. Otherwise, it is called "degenerate critical point".*

Lemma 2.1 *Let* p_0 *be a critical point of a smooth function* $f : M \to \mathbb{R}$*,* $(U, \varphi = (x_1, ..., x_n))$*,* $(V, \psi = (X_1, ..., X_n))$ *be two charts of* p_0*, and* $H_f(p_0), \mathcal{H}_f(p_0)$ *be the Hessians of* f *at* p_0*, using the charts* $(U, \varphi), (V, \psi)$ *respectively. Then the following holds:*

$$\mathcal{H}_f(p_0) = J(p_0)^t H_f(p_0) J(p_0) \tag{2.3}$$

where $J(p_0)$ *is the Jacobian matrix for the given coordinate transformation, defined by*

$$J(p_0) = \begin{bmatrix} \frac{\partial x_1}{\partial X_1}(p_0) & \cdots & \frac{\partial x_1}{\partial X_n}(p_0) \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial X_1}(p_0) & \cdots & \frac{\partial x_n}{\partial X_n}(p_0) \end{bmatrix} \tag{2.4}$$

and the matrix $J(p_0)^t$ *is the transpose of* $J(p_0)$*.*

For a critical point p_0, non-degeneracy does not depend on the choice of charts around p_0. The same argument is also true for degenerate critical points. In fact we have

$$\mathcal{H}_f(p_0) = J(p_0)^t H_f(p_0) J(p_0)$$

by the previous lemma,and hence

$$det\mathcal{H}_f(p_0) = det J(p_0)^t det H_f(p_0) det J(p_0) \tag{2.5}$$

by using determinant function on both sides. On the other hand, the determinant of the Jacobian matrix is non-zero. So the statement "$det\mathcal{H}_f(p_0) \neq 0$" and "$det H_f(p_0) \neq 0$" are equivalent. In other words,

$$det\mathcal{H}_f(p_0) \neq 0 \Leftrightarrow det H_f(p_0) \neq 0.$$

Now a function $f : M \to \mathbb{R}$ is called a *Morse function* if any critical point of f is non-degenerate. From now on, we only consider a Morse function f.

Now, we introduce Morse lemma on manifolds.

Theorem 2.4 *(The Morse Lemma) Let M be an n-dimensional smooth manifold and p_0 be a non-degenerate critical point of a Morse function $f : M \to \mathbb{R}$. Then, there exists a local coordinate system $(X_1, X_2, ..., X_n)$ around p_0 such that the coordinate representation of f has the following form:*

$$f = -X_1^2 - X_2^2 - ... - X_\lambda^2 + X_{\lambda+1}^2 + ... + X_n^2 + c \tag{2.6}$$

where $c = f(p_0)$ and p_0 corresponds to the origin $(0, 0, ..., 0)$.

One may refer to see [5] for the proof.

The number λ of minus signs in the equation (2.6) is the number of negative diagonal entries of the matrix $H_f(p_0)$ after diagonalization. By Sylvester's law, λ does not depend on how $H_f(p_0)$ is diagonalized. So, λ is determined by f and p_0. The number λ is called "the index of the non-degenerate critical point p_0". Obviously, λ is an integer between 0 and n.

Note that,

1. A non-degenerate critical point is isolated.
2. A Morse function on a compact manifold has only finitely many critical points [5].

3 A Morse function on $SO(n)$

In this section, we will define a Morse function on $SO(n)$.

The set of all $n \times n$ orthogonal matrices, $O(n) = \{A = (a_{ij}) \in M_n(\mathbb{R}) : AA^t = I_n\}$ is a group with matrix multiplication. From the definition of $O(n)$,

$$detA = \pm 1, \text{ for any } A \in O(n).$$

An orthogonal matrix with determinant 1 is called *rotation matrix* and the set of this kind of matrices is also a group, called *special orthogonal group* and denoted by $SO(n)$. On the other hand, let $S_n(\mathbb{R})$ denote the set of symmetric $n \times n$ matrices. Since each symmetric matrix is uniquely determined by its entries on and above the main diagonal, that is a linear subspace of $M_n(\mathbb{R})$ of dimension $n(n+1)/2$.

Now we define a function $\varphi : GL_n(\mathbb{R}) \longrightarrow S_n(\mathbb{R})$ by

$$\varphi(A) := A^t A.$$

Then, the identity matrix I_n is a regular value of φ [3].

Let $C \in S_n(\mathbb{R})$ with entries c_i, with $0 \leq c_1 < c_2 < ... < c_n$ fixed real numbers and $f_C : SO(n) \to \mathbb{R}$ be given by,

$$f_C(A) :=< C, A >= c_1 x_{11} + c_2 x_{22} + ... + c_n x_{nn}, \tag{3.1}$$

where $A = (x_{ij}) \in SO(n)$.

Obviously, f_C is a smooth function. Now, we will determine its critical points.

Lemma 3.1 *The critical points of the function f_C defined above are:*

$$\begin{bmatrix} \pm 1 & 0 & \cdots & 0 \\ 0 & \pm 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \pm 1 \end{bmatrix} \tag{3.2}$$

Proof.

Let A be a critical point of f_C. Then the derivative of f_C at A must be zero. Consider the matrix given by a rotation of first and second coordinate $B_{12}(\theta)$ defined by

$$B_{12}(\theta) = \begin{bmatrix} cos\theta & -sin\theta & 0 & \cdots & 0 \\ sin\theta & cos\theta & 0 & \cdots & 0 \\ 0 & 0 & 1 & & \vdots \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}.$$

Then, $AB_{12}(\theta) \in SO(n)$ and the matrix $B_{12}(\theta)$ forms a curve on $SO(n)$. Moreover, $B_{12}(\theta) = A$ for $\theta = 0$.

By the definition of f_C, and after computing the matrix product, we have

$$f_C(AB_{12}(\theta)) = c_1(x_{11}cos\theta + x_{12}sin\theta) + c_2(-x_{21}sin\theta + x_{22}cos\theta) + c_3x_{33} + \cdots + c_nx_{nn} \tag{3.3}$$

By differentiating f in the direction of the velocity vector $\frac{d}{d\theta}AB_{12}(\theta)|_{\theta=0}$ of the curve $AB_{12}(\theta)$ at A, we have

$$\frac{d}{d\theta} f_C(AB_{12}(\theta))|_{\theta=0} = c_1x_{12} - c_2x_{21} \tag{3.4}$$

and

$$\frac{d}{d\theta} f_C(B_{12}(\theta)A)|_{\theta=0} = -c_1x_{21} + c_2x_{12}. \tag{3.5}$$

However, by the assumption that A is a critical point of f_C, we require these derivatives to be zero. i.e.

$$c_1x_{12} - c_2x_{12} = 0$$

$$-c_1x_{21} + c_2x_{12} = 0$$

Solving this system for x_{12}, x_{21} gives $x_{12} = x_{21} = 0$. We can carry out the similar calculation for $B_{ij}(\theta)$ with $i < j$, where $B_{ij}(\theta)$ is with the entries: $(i,i) = cos\theta, (i,j) = -sin\theta, (j,i) = sin\theta$ and $(j,j) = cos\theta$. Thus, for the matrix A, $x_{ij} = 0$ whenever $i \neq j$. So, that is, a critical point of f_C is a diagonal matrix. On the other hand $A \in SO(n)$, so we have $AA^t = I_n$. So each entry on the main diagonal of A must be ± 1.

Conversely, let A be a matrix in the form (3.2). In order to check that A is a critical point, we need to compute the derivative of f_C. If we could find $n(n-1)/2$ curves C_i going through A with velocity vector at A and linearly independent from each other. Since the velocity vector of C_i at A plays a role of a local coordinate of A, we only need to check that the derivative of $f_C(C_i)$ vanishes to see that $Df(A) = 0$. Now, the claim is the curves C_i's are in fact $AB_{ij}\theta$' s defined above. Let $\epsilon_i = A_{ii}$ where A_{ii} is the i-th diagonal entry of $A(\epsilon_i = \pm 1)$. Then, the derivative of the matrix $AB_{ij}(\theta)$ at A is (we did for the case B_{12}, but it is same for other indices with $i < j$),

$$\frac{d}{d\theta} AB_{12}(\theta)|_{\theta=0} = \begin{bmatrix} 0 & -\epsilon_1 & 0 & \cdots & 0 \\ \epsilon_2 & 0 & 0 & \cdots & 0 \\ 0 & 0 & \ddots & & 0 \\ \vdots & \vdots & & & \vdots \\ 0 & 0 & \cdots & & 0 \end{bmatrix}$$

This matrix is regarded as a vector in $\mathbb{R}^{n^2}$. By considering all $1 \leq i \leq j \leq n$, these matrices (vectors) form a basis for the tangent space $T_A SO(n)$.

So, for a given matrix A in the form (3.2), it is easy to compute that, the derivative of f_C at A is zero. This means nothing but A is a critical point of f_C. ■

After now, we know the coordinate system of $SO(n)$ and the critical points of the the given function f_C. It is straightforward to compute the Hessian of f_C at A. Suppose that A is a critical matrix with diagonal entries $A_{ii} = \epsilon_i = \pm 1$. Then, we want to compute

$$\frac{\partial^2}{\partial\theta\partial\varphi} f_C(AB_{\alpha\beta}(\theta)B_{\gamma\delta}(\varphi))|_{\theta=0,\varphi=0}.$$

Notice that is linear $AB_{\alpha\beta}(\theta)B_{\gamma\delta}(\varphi)$ is linear in θ and in φ, and f_C is a linear function. Thus, we can bring the derivative inside f_C. So,

$$\begin{aligned} \frac{\partial^2}{\partial\theta\partial\varphi} f_C(AB_{\alpha\beta}(\theta)B_{\gamma\delta}(\varphi))|_{\theta=0,\varphi=0} &= f_C(A\frac{d}{d\theta}B_{\alpha\beta}(\theta)|_{\theta=0}\frac{d}{d\varphi}B_{\gamma\delta}(\varphi)|_{\varphi=0} \\ &= \begin{cases} -c_\alpha\epsilon_\alpha - c_\beta\epsilon_\beta & \text{if } \alpha = \gamma,\ \beta = \delta \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

This calculation becomes easier if we consider the matrix multiplication $c_{ij} = \sum_k a_{ik}b_{kj}$. The calculation above shows that the Hessian matrix is diagonal. Since $c_\alpha \neq c_\beta$ for $\alpha \neq \beta$, the entries on the diagonal is non-zero. Therefore, A is a non-degenerate critical point of f_C, meaning that f_C is a Morse function on $SO(n)$.

Assume that the subscripts i of the diagonal entries ϵ_i of A , $1 \leq i \leq n$, with $\epsilon_i = 1$ are

$$i_1, i_2, \cdots, i_m$$

in ascending order. Then the index of the critical point A (the number of minus signs on the diagonal of Hessian) is

$$(i_1 - 1) + (i_2 - 1) + \cdots + (i_m - 1).$$

And the index is 0 if all ϵ_i' s are -1. Also, the critical value at the critical point is

$$2(c_{i1} + c_{i2} + \cdots + c_{im}) - \sum_{i=0}^{n} c_i.$$

Considering that $detA = 1$, there are 2^{n-1} critical points [4]

4 Perfect Morse Functions

First, we will give the basic notions.

Definition 4.1 *The Poincaré polynomial of the* n*- dimensional manifold* M *is defined to be*

$$P_M(t) = \sum_{k=0}^{n} b_k(M) t^k \tag{4.1}$$

where $b_k(M)$ *is the* k*-th Betti number of* M*.*

Definition 4.2 *Let* $f : M \to \mathbb{R}$ *be a Morse function. Then, the Morse polynomial of* f *is defined to be*

$$P_f(t) = \sum_{k=0}^{n} \mu_k t^k \tag{4.2}$$

where μ_k *is the number of critical points of* f *of index* k*.*

Theorem 4.3 *(The Morse Inequality) Let* $f : M \to \mathbb{R}$ *be a Morse function on a smooth manifold* M*. Then, there exists a polynomial* $R(t)$ *with non-negative integer coefficients such that*

$$P_f(t) = P_M(t) + (1 + t)R(t).$$

One may refer to [7] for proof.

A Morse function $f : M \to \mathbb{R}$ is called a perfect Morse function if $P_f(t) = P_M(t)$ [7].

Now, we show that the function f_C on $SO(n)$ defined in the previous section is also a perfect Morse function.

Theorem 4.4 *The function*

$$f_C : SO(n) \to \mathbb{R}, \quad f_C(A) := \langle C, A \rangle$$

is a perfect Morse function, where $C \in S_n(\mathbb{R})$*.*

Proof. First we show that the Morse polynomial is,

$$P_{f_C}(t) = (1+t)(1+t^2)\cdots(1+t^{n-1}). \tag{4.3}$$

We use induction method. For making it easier, we label the function f_C with n as $f_{C_n} : SO(n) \to \mathbb{R}$.
Trivially, for $n = 1$, $P_{f_{C_1}}(t) = 1$ and for $n = 2$, $P_{f_{C_2}}(t) = 1 + t$. Assume that, $P_{f_{C_n}}(t) = (1+t)(1+t^2)+\cdots+(1+t^{n-1})$. Then, we need to show that $P_{f_{C_{n+1}}}(t)$ satisfies the form (4.3).
We may consider that $SO(n+1)$ gets all the critical points from $SO(n)$ with extra bottom entry ($(n+1)$- th diagonal entry), which is either +1 or -1. Say the set of all these points are C^+_{n+1} and C^-_{n+1} respectively.
Let $A \in C^-_{n+1}$. Then we have $\tilde{A} \in O(n)$ such that, A is the matrix $\tilde{A}$ with extra bottom entry -1. Then, by the definition of index, we obtain

$$ind(A) = ind(\tilde{A}).$$

Thus, for the elements of C^-_{n+1} the equation (4.3) holds. Let $A \in C^+_{n+1}$. Then we have $\tilde{A} \in SO(n)$ such that, A is the matrix with $\tilde{A}$ with the bottom entry +1. Thus, by the definition of index, we obtain

$$ind(A) = ind(\tilde{A}) + n.$$

So, by the definition of Morse polynomial, we gain

$$P_{f_{C_{n+1}}}(t) = P_{f_{C_n}}(t)(1+t^n) = (1+t)(1+t^2)\cdots(1+t^{n-1})(1+t^n) \tag{4.4}$$

.

Now, we find out the Poincaré polynomial of $SO(n)$. The graded abelian group $H_*(SO(n), \mathbb{Z}_2)$ is isomorphic to the graded group coming from the exterior algebra

$$\wedge_{\mathbb{Z}_2}[e_1, e_2, ..., e_{n-1}]$$

[2]. Let say $A(n) = \wedge_{\mathbb{Z}_2}[e_1, e_2, ..., e_{n-1}]$ where the degree of e_i, $|e_i| = i$. Then, we obtain

$$|e_{i_1} \wedge e_{i_2} \wedge ... \wedge e_{i_k}| = \sum_{j=1}^{k} |e_{i_j}| = \sum_{j=1}^{k} i_j.$$

If we define $a(n)_k = dim\mathbb{Z}_2(A(n)_k)$, then by the result in [2], $a(n)_k$ is nothing but the k- th Betti number of $SO(n)$. Hence, the polynomial

$$P(A(n)) = \sum_{i=0}^{\infty} a(n)_i t^i$$

is the Poincaré polynomial of $SO(n)$.

Now, our claim is that The Poincaré polynomial of $SO(n)$ is

$$P(A(n)) = (1+t)(1+t^2)\cdots(1+t^{n-1}).$$

Let $B(A(n))$ be the basis of $A(n)$. For instance, $B(A(1)) = trivial$, $B(A(2)) = \{1, e_1\}$, $B(A(3)) = \{1, e_1, e_2, e_1 \wedge e_2\}$ etc.

In this sense, we obtain

$$B(A(n+1)) = (B(A(n)) \wedge e_n) \sqcup B(A(n)).$$

We use induction method. Indeed, here we have very similar arguments with the previous claim. The variable e_n has the same role with " the extra bottom entry ±1". Then, we have the polynomial $P(A(n)) = \sum_{b \in B(A(n))} a(n)_b t^{|b|}$. Trivially, $P(A(1)) = 1$ and $P(A(2)) = 1 + t$. By the induction hypothesis, assume that

$$P(A(n)) = \sum_{b \in B(A(n))} a(n)_b t^{|b|} = (1+t)(1+t^2)\cdots(1+t^{n-1}).$$

For the polynomial $P(A(n+1))$, pick an element $b \in B(A(n+1))$. Then, b is in either $B(A(n))$ or $B(A(n)) \wedge e_n$. For $b \in B(A(n+1))$, trivially, $P(A(n+1))$ has the desired form. If $b \in B(A(n)) \wedge e_n$, then by the definition of degree, there is $\tilde{b} \in B(A(n))$ such that $|b| = |\tilde{b}| + n$. Thus, by the definition of $P(A(n))$, we obtain

$$P(A(n+1)) = (1+t)(1+t^2)\cdots(1+t^n) \tag{4.5}$$

which completes the proof. ■

Thereby, we have shown that, for the given Morse function $f_C : SO(n) \to \mathbb{R}$, $P_M(t) = P_{f_C}(t)$, meaning that f_C is a perfect Morse function.

References

[1] D. B. Gauld. *Differential topology: An Introduction.* Marcel Dekker, New York, 1982.

[2] A. Hatcher. *Algebraic Topology.* Cambridge University Press, New York, 3rd edition, 2002.

[3] J. M. Lee. *Introduction to Smooth Manifolds.* Springer, New York, 2rd edition, 2012.

[4] Y. Matsumoto. *An Introduction to Morse Theory.* American Mathematical Society, 2002.

[5] J. W. Milnor. *Morse Theory.* Princeton University Press, 1963.

[6] M. Morse. The foundations of a theory of the calculus of variations in the large in m-space. *Trans. Amer. Math. Soc.*, 30:213–274, 1928.

[7] L. I. Nicolaescu. *An Invitation to Morse Theory.* Springer, New York, 2rd edition, 2011.